



Variations of crustal elastic properties during the 2009 L'Aquila earthquake inferred from cross-correlations of ambient seismic noise.

Lucia Zaccarelli, N.M. Shapiro, L. Faenza, G. Soldati, Alberto Michelini

► To cite this version:

Lucia Zaccarelli, N.M. Shapiro, L. Faenza, G. Soldati, Alberto Michelini. Variations of crustal elastic properties during the 2009 L'Aquila earthquake inferred from cross-correlations of ambient seismic noise.. Geophysical Research Letters, 2011, VOL. 38, pp.6 PP. 10.1029/2011GL049750 . hal-00706545

HAL Id: hal-00706545

<https://hal.science/hal-00706545>

Submitted on 11 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Variations of crustal elastic properties during the 2009
2 L'Aquila earthquake inferred from cross-correlations of
3 ambient seismic noise

L. Zaccarelli¹, N. M. Shapiro¹, L. Faenza², G. Soldati², A. Michelini²

L. Zaccarelli, Institut de Physique du Globe de Paris, Sorbonne Paris Cité, CNRS (UMR7154),
1 rue Jussieu, 75238 Paris, cedex 5, France. (zaccarelli@ipgp.fr)

N.M. Shapiro, Institut de Physique du Globe de Paris, Sorbonne Paris Cité, CNRS (UMR7154),
1 rue Jussieu, 75238 Paris, cedex 5, France. (nshapiro@ipgp.fr)

L. Faenza, Istituto Nazionale di Geofisica e Vulcanologia, via di Vigna Murata 605, 00143
Roma, Italy. (licia.faenza@ingv.it)

G. Soldati, Istituto Nazionale di Geofisica e Vulcanologia, via di Vigna Murata 605, 00143
Roma, Italy. (gaia.soldati@ingv.it)

A. Michelini, Istituto Nazionale di Geofisica e Vulcanologia, via di Vigna Murata 605, 00143
Roma, Italy. (alberto.michelini@ingv.it)

¹Institut de Physique du Globe de Paris,

4 We retrieve seismic velocity variations within the Earth's crust in the re-
5 gion of L'Aquila (central Italy) by analyzing cross-correlations of more than
6 two years of continuous seismic records. The studied period includes the April
7 6, 2009, M_W 6.1 L'Aquila earthquake. We observe a decrease of seismic ve-
8 locities as a result of the earthquake's main shock. After performing the anal-
9 ysis in different frequency bands between 0.1 and 1 Hz, we conclude that the
10 velocity variations are strongest at relatively high frequencies (0.5-1 Hz) sug-
11 gesting that they are mostly related to the damage in the shallow soft lay-
12 ers resulting from the co-seismic shaking.

Sorbonne Paris Cité, CNRS (UMR7154),

France.

²Istituto Nazionale di Geofisica e

Vulcanologia, Roma, Italy.

1. Introduction

On April 6, 2009 a M_W 6.1 earthquake struck the central Apennines region near L'Aquila (Italy) causing severe damage and more than 300 fatalities [Scognamiglio *et al.*, 2010]. This area had been long recognized as seismically active [see the official seismic hazard map of Italy, *MPS Working Group*, 2004] and an occurrence of a strong earthquake in the central Apennines could not be considered as totally unexpected. Before the main shock, an increase in the rate of seismicity started on September 2008 and many small size events (about 300 with $M_L \leq 2.5$) occurred beneath the L'Aquila city area. This foreshock sequence culminated with a $M_L = 4.1$ earthquake on March 30, 2009. In the following days, the seismicity decreased until two earthquakes ($M_L = 3.9$ and $M_L = 3.5$) occurred just a few hours before the L'Aquila main shock. In agreement with the extensional tectonics of the central Apennines, the focal mechanism of the L'Aquila earthquake has been determined to be a normal fault on a South-West dipping plane with the rupture area of $\sim 20 \times 15$ km² and the dipping angle of about 50 degrees [Cirella *et al.*, 2009]. The main shock was followed by an aftershock sequence that included 33 earthquakes greater than $M_L = 4$.

In this study, we use a recently proposed monitoring technique based on ambient seismic noise. The idea of this method is to use signals reconstructed from repeated cross-correlations of continuous seismic records as virtual seismograms generated by highly repeatable sources. In case of well distributed noise, the reconstructed virtual sources are close to point forces and the cross-correlations functions can be considered as Green functions [e.g., Weaver and Lobkis, 2001; Shapiro and Campillo, 2004; Sabra *et al.*, 2005;

Shapiro *et al.*, 2005]. Highly accurate temporal monitoring can be also performed even with inhomogeneous noise sources distributions when a perfect reconstruction of the Green function is not achieved [e.g., Hadziioannou *et al.*, 2009]. The changes of the travel times measured from the noise cross-correlations reflect variations of the elastic properties in the propagating media, i.e., in the Earth's crust. This approach has been recently applied to monitor active volcanoes [e.g., Sens-Schönfelder and Wegler, 2006; Brenguier *et al.*, 2008a; Duputel *et al.*, 2009; Mordret *et al.*, 2010] and large seismogenic faults [e.g., Wegler and Sens-Schönfelder, 2007; Brenguier *et al.*, 2008b; Chen *et al.*, 2010] and to detect seasonal changes in the Earth's crust resulting from thermoelastic variations [e.g., Meier *et al.*, 2010].

In a seismogram or in a correlation function, the delay accumulates linearly with the lapse time when the wave speed changes homogeneously within the medium. As a consequence, a small change can be detected more easily when considering late arrivals. This makes the use of coda waves particularly suited to measure temporal variations. This can be done either by using the so-called stretching technique [e.g., Wegler and Sens-Schönfelder, 2007] or with a method that was initially developed for repeated earthquakes (doublets) by Poupinet *et al.* [1984]. Here, we use this latter approach that has been specifically adopted to make measurements from the noise cross-correlations [e.g., Clarke *et al.*, 2011]. We apply this method to two years of continuous recordings by three seismic stations located in the vicinity of the L'Aquila main shock fault (Figure 1) to measure variations of crustal seismic velocities caused by this earthquake.

2. Selecting and pre-processing the data and computing cross-correlations

Istituto Nazionale di Geofisica e Vulcanologia (INGV) operates two large seismological networks: the Italian National Seismic Network (INSN) and the Mediterranean Very Broadband Seismographic Network MedNet. The INSN consists of more than 250 stations with various characteristics [Amato and Mele, 2008]. MedNet consists of 22 very broadband stations distributed over the Euro-Mediterranean area with 13 of them located in Italy [Mazza *et al.*, 2008]. During period of interest for our study, four broadband stations operated in continuous mode within a radius of 25 km from the main shock epicenter. However, records of one of these stations contained too many gaps and we finally decided to use three stations: CAMP and FIAM from INSN and AQU from MedNet (Figure 1). The longest period of data availability at these three stations was between March 27, 2008 and April 18, 2010.

We re-sampled time series recorded at the three stations in order to get a perfect time synchronization and filled existing small gaps via a linear interpolation. Then, we pre-processed the vertical component seismograms by whitening their spectra between 0.1 and 1 Hz and by normalizing their amplitude through a one-bit normalization. In this way, the contributions arising from strong transient phenomena were reduced in both time and frequency domains [e.g., Bensen *et al.*, 2005; Brenguier *et al.*, 2008b]. Finally, we computed cross-correlations between the three pairs of stations for every hour of the available recordings.

3. Measurement of seismic velocity variations

We adopted the Multi Window Cross-Spectrum (MWCS) analysis [e.g., *Clarke et al.*, 2011]. This technique was first proposed by *Poupinet et al.* [1984] for retrieving the relative velocity variation between earthquake doublets. *Brenguier et al.* [2008a, b] applied this technique to the cross-correlations of the seismic noise. The main idea of the method is that the noise cross-correlations computed from subsequent time windows can be analysed similar to records from earthquake doublets. When analyzing long time series, we compare a single reference cross-correlation with many subsequent current functions. The reference cross-correlation CC^R for a particular station pair is obtained from stacking all available cross-correlations for this pair and, therefore, is representative of the background crustal state. The current cross-correlations CC^C are obtained from stacking a small sub-set of cross-correlations representative of a state of the crust for a given short period of time. There is a trade-off between the length of the stack required to have stable estimates of the CC^C and the time resolution for detecting the variations. To find an optimal stacking duration for the current function we tested different lengths between 10 and 100 days. For each tested stacking length, we computed all possible functions CC^C by applying moving windows shifted by two days. Then, we computed the correlation coefficient r between the reference function CC^R and every CC^C . The distribution of r characterizes the similarity between CC^R and CC^C for a given stacking length. We represent the overall degree of similarity by the mean and the standard deviation of this distribution. Figure 2 shows these parameters for the three station pairs. We observe that the degree of similarity

94 increases rapidly for short stacking durations and then it tends to stabilize. We selected
 95 a value of 50 days as stacking length for computing the current correlation functions.

96 The MWCS analysis consists of two computational steps [e.g., *Clarke et al.*, 2011]. In
 97 the first step, we estimate for a station pair k delay times δt_i^k between CC^R and CC^C
 98 within a set of time windows centered at t_i . In case of uniform velocity perturbations,
 99 the measured delays δt_i^k are expected to be a linear function of time t_i with a slope
 100 corresponding to the relative time perturbation:

$$\frac{\Delta t}{t} = -\frac{\Delta v}{v} \quad (1)$$

101 where $\frac{\Delta v}{v}$ is the relative uniform seismic velocity perturbation that can be estimated in
 102 the second step from a single station pair k via linear fitting of the following equation:

$$\delta t_i^k = -\left(\frac{\Delta v}{v}\right)_k \cdot t_i \quad (2)$$

103 In order to obtain one estimates representative of the entire crustal volume, we merged
 104 together the delays δt_i^k measured from the three station pairs before proceeding with the
 105 second step of the analysis. We computed the median value $\widetilde{\delta t_i}$ of the delays δt_i^k for every i -
 106 th window, and we inserted it into (2) to estimate of $\frac{\Delta v}{v}$ for the entire region encompassed
 107 by the three stations. When performing this analysis, we removed the central part of
 108 the cross-correlations containing direct waves (group velocities faster than 2.5 km/s; see
 109 Table 1) because they may be sensitive to the changing position of the noise sources [e.g.,
 110 *Froment et al.*, 2010]. Relative velocity variations were then computed by taking into

account the coda of the cross-correlation up to a length of 60 s where the signal decreases to values close to the noise level.

To estimate uncertainties of our measurements, we followed the method proposed by *Clarke et al.* [2011] and performed a synthetic test on the L'Aquila noise cross-correlations. We perturbed the reference cross-correlation function by stretching its waveform and simulating different values of velocity variations (from 0.01% to 0.5%). Then, we added a random noise with a signal-to-noise ratio of 5 (that is the mean value measured from the observed cross-correlations). Finally, we applied the MWCS technique to measure the apparent velocity variations $\frac{\Delta v}{v}$ between the perturbed cross-correlations and the original CC^R . The RMS deviations between the estimated velocity variations and those introduced through stretching characterize the uncertainties of our measurements.

To investigate the depth extent of the measured crustal velocity perturbations, we performed the MWCS analysis inside three different frequency bands: [0.1–1], [0.1–0.6], and [0.5–1] Hz. It has been shown both theoretically and observationally that at these frequencies the coda of seismograms and correlation functions are mainly composed of surface waves [e.g., *Hennino et al.*, 2001; *Margerin et al.*, 2009]. We therefore expect that the sensitivity of the coda waves to a velocity change at depth depends on their spectral content with shorter periods sensitive to shallower structures and longer periods sampling deeper parts of the crust. The measurement results for the three frequency bands are presented in **Figure 3** and show a sudden velocity decrease at the time of occurrence of the L'Aquila main shock. The amplitude of this velocity drop is largest at frequencies

132 higher than 0.5 Hz and decreases at lower frequencies. This indicates that a large part of
 133 the observed variations have their likely origin within the shallow crustal layers.

4. Discussion

134 A limited number of available stations (only three) and the fact that only
 135 one of them is located in the immediate vicinity of the earthquake fault did
 136 not allow us to identify exact regions that produced the observed velocity
 137 variations. Also, the dataset used in this study did not allow us to make
 138 robust measurements with refined time resolution. A denser network covering
 139 the source area would be required to obtain better space and time resolutions
 140 [e.g., *Brenquier et al.*, 2008a]. Therefore, we interpret here only the most
 141 robust average features.

142 The results presented in our study show that the L'Aquila main shock caused a de-
 143 tectable reduction of seismic velocities within the surrounding crust. We observe that
 144 the velocity dropped by 0.3%, which is more than 3 times larger than the fluctuations
 145 observed before the main shock. Co-seismic velocity reductions can be attributed to
 146 increasing crack and void densities in the shallow crustal structure and/or to reduced
 147 compaction of the near-surface granular material. The presence and migration of fluids
 148 can further contribute to modification of the seismic properties in the shallow crust. Our
 149 results can be compared with other studies that have addressed changes of the crustal
 150 parameters prior and after the L'Aquila earthquake. *Amoruso and Crescentini* [2010]
 151 used strain measurements obtained in the Gran Sasso laboratory during the two years
 152 prior to the main shock to infer that no anomalous signal was observed. They concluded

that the possible earthquake nucleation zone was confined to a volume less than 100 km^3 . In contrast, v_p/v_s anomalies have been reported by *Di Luccio et al.* [2010] in the weeks prior to the main shock with an abrupt variation after the $M_L = 4.1$ foreshock occurred on March 30. Similar results were obtained by *Lucente et al.* [2010] who used shear wave splitting in addition to v_p/v_s ratios. They attribute the velocity anomalies occurring in the week prior to the main shock to a complex sequence of dilatancy-diffusion processes in which fluids play a key role. *Terakawa et al.* [2010] inverted the stress field obtained from the aftershock sequence focal mechanisms to determine the fluid pressure and to conclude that the spatial pattern of the sequence is driven mainly by fluid migration.

Our results are based on current cross-correlation functions stacked over a 50 day period and, therefore, do not have the time resolution required to identify possible short-term precursory variations and to separate them from the co-seismic effect. On the other hand, with stacking large data volumes our estimation of the co-seismic velocity reduction is inherently very robust. The observed velocity reduction is larger at higher frequencies. Therefore, we prefer the hypothesis the perturbation is mainly due to damaging of shallow soft sedimentary layers by the co-seismic strong ground motion [e.g., *Wu et al.*, 2009]. This effect may be also enhanced by the presence of fluids.

We compare the co-seismic perturbation observed during the L'Aquila earthquakes with other cases when the co-seismic crustal velocity variations were measured from noise cross-correlations (Table 2). The co-seismic velocity drop measured for the L'Aquila earthquake ($\sim 0.3\%$) is significantly larger than the values measured within a similar frequency band for the M_W 6.0 Parkfield and the M_W 7.9 Wenchuan events ($\Delta v/v \sim 0.08\%$ as reported by

175 *Brenquier et al.* [2008a] and *Chen et al.* [2010], respectively). At the same time, a stronger
 176 variation ($\sim 0.6\%$) has been observed **with** the stretching technique and frequencies higher
 177 than 2 Hz during the M_W 6.6 Mid-Niigata earthquake. The results of this comparison
 178 suggest that the level of measured co-seismic velocity variation is not a simple function of
 179 the total moment release during an earthquake but is controlled by different factors such
 180 as local geological conditions and, possibly, focal mechanism and source depth. Also, the
 181 frequency range used in the analysis controls the depth extent of the measured anomaly.
 182 Finally, the aperture of the used seismic network (i.e., the distance between the station
 183 pairs) can play an important role. So far, the velocity variations reported in this study
 184 were measured over a relatively large area. Therefore, they may be less sensitive to the
 185 processes occurring in the immediate vicinity of the fault, where stress-induced velocity
 186 perturbations are expected to be most important.

187 **Acknowledgments.** The data used in this study were provided by the Istituto
 188 Nazionale di Geofisica e Vulcanologia. We thank G. Moguilny for maintaining the Cohersis
 189 cluster on which all computations were performed. This work was supported by Agence
 190 Nationale de la Recherche (France) under contract ANR-06-CEXC-005 (COHERSIS) and
 191 by a FP7 ERC Advanced grant 227507 (WHISPER).

References

192 Amato, A., and F. Mele, (2008), Performance of the INGV National Seismic Network
 193 from 1997 to 2007, *Annals of Geophysics*, 51, 417-431.

- 194 Amoruso, A., and L. Crescentini, (2010), Limits on earthquake nucleation and other
195 pre-seismic phenomena from continuous strain in the near field of the 2009 L'Aquila
196 earthquake, *Geophys. Res. Lett.*, *37*, L10307, doi:10.1029/2010GL043308.
- 197 Bensen, G.D., M.H. Ritzwoller, M.P. Barmin, A.L. Levshin, F. Lin, M.P. Moschetti, N.M.
198 Shapiro, and Y. Yang (2007), Processing seismic ambient noise data to obtain reliable
199 broad-band surface wave dispersion measurements, *Geophys. J. Int.*, *169*, 1239-1260,
200 doi: 10.1111/j.1365-246X.2007.03374.x.
- 201 Brenguier, F., M. Campillo, C. Hadziioannou, N. M. Shapiro, R. M. Nadeau, and E.
202 Larose (2008a), Postseismic relaxation along the San Andreas fault at Parkfield from
203 continuous seismological observations, *Science*, *321*, 1478–1481.
- 204 Brenguier, F., N. M. Shapiro, M. Campillo, V. Ferrazzini, Z. Duputel, O. Coutant, and A.
205 Nercessian (2008b), Towards forecasting volcanic eruptions using seismic noise, *Nature*
206 *Geoscience*, *1*, 126–130.
- 207 Chen, J. H., B. Froment, Q. Y. Liu, and M. Campillo (2010), Distribution of seismic
208 wave speed changes associated with the 12 May 2008 Mw 7.9 Wenchuan earthquake,
209 *Geophys. Res. Lett.*, *37*, L18302, doi:10.1029/2010GL044582.
- 210 Chiarabba, C., and 29 others (2009), The 2009 L'Aquila (central Italy) M_W
211 6.3 earthquake: Main shock and aftershocks, *Geophys. Res. Lett.*, *36*, L18308,
212 doi:10.1029/2009GL039627.
- 213 Cirella A., A. Piatanesi, M. Cocco, E. Tinti, L. Scognamiglio, A. Michelini, A. Lomax, and
214 E. Boschi (2009), Rupture history of the 2009 L'Aquila (Italy) earthquake from non-
215 linear joint inversion of strong motion and GPS data, *Geophys. Res. Lett.*, *36*, L19304,

doi:10.1029/2009GL039795.

Clarke, D., L. Zaccarelli, N.M. Shapiro, and F. Brenguier (2011), Assessment of resolution and accuracy of the Moving Window Cross Spectral technique for monitoring crustal temporal variations using ambient seismic noise, *Geophys. J. Int.*, *186*, 867-882, doi: 10.1111/j.1365-246X.2011.05074.x.

Di Luccio, F., G. Ventura, R. Di Giovambattista, A. Piscini, and F. R. Cinti (2010), Normal faults and thrusts re-activated by deep fluids: the 6 April 2009 M_W 6.3 L'Aquila earthquake, central Italy, *J. Geophys. Res.*, *115*, B06315, doi: 10.1029/200JB007190.

Duputel, Z., V. Ferrazzini, F. Brenguier, N. Shapiro, M. Campillo, and A. Nercessian (2009), Real time monitoring of relative velocity changes using ambient seismic noise at the Piton de la Fournaise volcano (La Reunion) from January 2006 to June 2007, *J. Volcanol. Geotherm. Res.*, *184*, 164-173, doi:10.1016/j.jvolgeores.2008.11.024.

EMERGEO Working Group (2010), Evidence for surface rupture associated with the M_W 6.3 L'Aquila earthquake sequence of April 2009 (central Italy), *Terra Nova*, *22*, 43-51.

Froment, B., M. Campillo, P. Roux, P. Gouedard, A. Verdel, and R.L. Weaver (2010), Estimation of the effect of nonisotropically distributed energy on the apparent arrival time in correlations, *Geophysics*, *75*, SA85-SA93.

Hadziioannou, C., E. Larose, O. Coutant, P. Roux, and M. Campillo (2009), Stability of Monitoring Weak Changes in Multiply Scattering Media with Ambient Noise Correlation: Laboratory Experiments, *J. Acoust. Soc. Am.*, *125*, 3688-3695.

Hennino R., N. Trégoures, N. M. Shapiro, L. Margerin, M. Campillo, B.A. Van Tiggelen, and R.L. Weaver (2001), Observation of Equipartition of Seismic Waves, *Phys. Rev. Lett.*,

86, 3447–3450.

Lucente, F. P., P. De Gori, L. Margheriti, D. Piccinini, M. Di Bona, C. Chiarabba, and N. Piana Agostinetti (2010), Temporal variations of seismic velocity and anisotropy before the 2009 M_w 6.3 L'Aquila earthquake, Italy, *Geology*, *38*, 11, 1015–1018, doi:10.1130/G31463.1.

Margerin L., M. Campillo, B. A. Van Tiggelen, and R. Hennino (2009), Energy partition of seismic coda waves in layered media: theory and application to Pinyon Flats Observatory, *Geophys. J. Int.*, *177*, 571–585.

Mazza S., M. Olivieri, A. Mandiello and P. Casale (2008), The Mediterranean Broad Band Seismographic Network Anno 2005/06, *Earthquake Monitoring and Seismic Hazard Mitigation in Balkan Countries, NATO Science Series*, *81*, 133–149, DOI: 10.1007/978-1-4020-6815.

Meier, U., N.M. Shapiro, and F. Brenguier (2010), Detecting seasonal variations in seismic velocities within Los Angeles basin from correlations of ambient seismic noise, *Geophys. J. Int.*, *181*, 985–996, doi: 10.1111/j.1365-246X.2010.04550.x.

Mordret, A., A.D. Jolly, Z. Duputel, and N. Fournier (2010), Monitoring of phreatic eruptions using Interferometry on Retrieved Cross-Correlation Function from Ambient Seismic Noise: Results from Mt. Ruapehu, New Zealand, *Journal of Volcanology and Geothermal Research*, *191*, 46–59.

MPS Working Group (2004). Redazione della mappa di pericolosità sismica prevista dall'Ordinanza PCM 3274 del 20 Marzo 2003, Rapporto Conclusivo per il Dipartimento della Protezione Civile, [http:// zonesismiche.mi.ingv.it](http://zonesismiche.mi.ingv.it) (last accessed: June

- 260 2011), INGV, Milano-Roma, 2004 April, 65 pp. including 5 appendixes.
- 261 Poupinet, G., W. L. Ellsworth, and J. Frechet (1984), Monitoring velocity variations in
262 the crust using earthquake doublets: an application to the Calaveras Fault, California,
263 *J. Geophys. Res.*, *89*, 5719–5731.
- 264 Roux, P., K. G. Sabra, P. Gerstoft, and W. A. Kuperman (2005), P-
265 waves from cross-correlation of seismic noise, *Geophys. Res. Lett.*, *32*, L19303,
266 doi:10.1029/2005GL023803.
- 267 Sabra, K. G., Gerstoft, P., Roux, P., Kuperman, W. A., and Fehler, M. C. (2005), Ex-
268 tracting time domain Green's function estimates from ambient seismic noise, *Geophys.*
269 *Res. Lett.*, *32*, L03310.
- 270 Scognamiglio L., E. Tinti, A. Michelini, A. S. Dreger, A. Cirella, M. Cocco, S. Mazza, A.
271 Piatanesi (2010), Fast determination of moment tensors and rupture history: what has
272 been learned from the 6 april 2009 L'Aquila earthquake sequence, *Seismol. Res. Lett.*,
273 *81* (6), pp. 892-906, doi:10.1785/gssrl.81.6.892.
- 274 Sens-Schönfelder, C. and U. Wegler (2006), Passive image interferometry and seasonal
275 variations of seismic velocities at Merapi Volcano, Indonesia, *Geophys. Res. Lett.*, *33*,
276 L21302, doi:10.1029/2006GL027797.
- 277 Shapiro, N. M., and M. Campillo (2004), Emergence of broadband Rayleigh waves
278 from correlations of the ambient seismic noise, *Geophys. Res. Lett.*, *31*, L07614,
279 doi:10.1029/2004GL019491.
- 280 Shapiro, N.M, M. Campillo, L. Stehly, and M.H. Ritzwoller (2005), High resolution surface
281 wave tomography from ambient seismic noise, *Science*, *307*, 1615-1618.

- 282 Terakawa, T., A. Zoporowski, B. Galvan, and S. Miller, (2010), High-pressure fluid at
283 hypocentral depths in the L'Aquila region inferred from earthquake focal mechanism,
284 *Geology*, *38*, 11, 995–998, doi:10.1130/G31457.1.
- 285 VanDecar, J. C., and R. S. Crosson, (1990) Determination of teleseismic relative phase
286 arrival times using multi-channel cross-correlation and least squares, *Bull. Seismol. Soc.*
287 *Am.*, *80*, 1, 150–169.
- 288 Weaver, R. L. and O. I. Lobkis (2001), Ultrasonics without a source: ther-
289 mal fluctuation correlations at MHz frequencies, *Phys. Rev. Lett.*, *87*, 134301,
290 doi:10.1103/PhysRevLett.87.134301.
- 291 Wegler, U., Sens-Schönfelder C. (2007), Fault zone monitoring with passive image inter-
292 ferometry, *Geophys. J. Int.*, *168*, 1029–1033.
- 293 Wu, C., Z. Peng, and Y. Ben-Zion (2009), Non-linearity and Temporal Changes of Fault
294 Zone Site Response Associated with Strong Ground Motion, *Geophys. J. Int.*, *176*,
295 265–278, doi: 10.1111/j.1365-246X.2008.04005.x.

Table 1. Parameters of the three inter-stations paths used in the study. The Rayleigh wave arrival times are roughly estimated considering a group velocity of 3 km/s [*Chiarabba et al.*, 2009]). Parts of the correlation functions with group velocities faster than 2.5 km/s we excluded from the analysis to avoid the influence of the noise source variability in direct arrivals.

| stations | distance km | Rayleigh arrival s | cutoff s |
|-----------|----------------|-----------------------|-------------|
| AQU_CAMP | 20 | 6.67 | ± 7.5 |
| AQU_FIAM | 26 | 8.67 | ± 10 |
| CAMP_FIAM | 38 | 12.67 | ± 15 |

Table 2. Comparison between the L'Aquila event and other earthquakes where co-seismic velocity variations were measured from noise cross-correlations. Values of velocity variations are from *Brenguier et al.* [2008a], *Wegler and Sens-Schönfelder* [2007], and *Chen et al.* [2010], for the Parkfield, the Mid-Niigata, and the Wenchuan earthquakes, respectively.

| Earthquake | M_w | depth km | focal mechanism | $\Delta v/v$ % | frequency Hz | stations |
|-------------|-------|-------------|-----------------|--------------------|---------------------------|----------|
| L'Aquila | 6.1 | 8.8 | normal | 0.15 0.3 0.4 | 0.1–0.6 0.1–1 0.5–1 | 3 |
| Parkfield | 6.0 | 7.9 | strike-slip | 0.08 | 0.1–0.9 | 13 |
| Mid-Niigata | 6.6 | 5 | thrust | 0.6 | > 2 | 1 |
| Wenchuan | 7.9 | 19 | thrust | 0.08 | 0.3–1 | > 30 |




Figure 1. Map of the central Apennines showing the location of the L'Aquila epicenter (black open star) and of the fault plane projection (black rectangle) from *Cirella et al.* [2009]. The gray triangles are the three stations considered in this study. Black thin lines indicate main tectonic faults from *Emergeo Working Group* [2010]. Light gray lines show the regional boundaries.




Figure 2. Mean (a) and standard deviation (b) values of the correlation coefficients r between CC^C and CC^R as a function of number of days used to construct the current correlation functions CC^C . Mean and standard deviations were computed after a Fisher transformation that returns an almost normally distributed variable [VanDecar and Crosson, 1990]. Panels (c), (d), and (e) show the reference cross-correlation functions CC^R (blue) together with an example of 50 day current function CC^C (black) for the three couples of stations. Only portions of the the signal considered in the analysis are plotted (Table 1). Numbers in the bottom left corners are the respective correlation coefficients r .




Figure 3. Relative velocity variations measured from cross-correlations of seismic noise recorded at the three stations (gaps correspond to periods when the stations were not operating simultaneously). Results obtained by analyzing the whole frequency range [0.1 1] Hz are shown with a gray color. Blue color shows the results from narrower frequency ranges: (a) [0.1 0.6] Hz and (b) [0.5 1] Hz). Vertical bars indicate the uncertainties of the measurements. The vertical red line highlights the time of occurrence of the L'Aquila main shock.